



## REVIEW ARTICLE

# The diagnostic and prognostic value of red cell distribution width in cardiovascular disease; current status and prospective

Seyed M. Parizadeh<sup>1</sup> | Reza Jafarzadeh-Esfehani<sup>2</sup> | Amirhossein Bahreyni<sup>3</sup> |  
Maryam Ghandehari<sup>4</sup> | Mojtaba Shafiee<sup>4,5</sup> | Farzad Rahmani<sup>6</sup> |  
Mohammar R. Parizadeh<sup>1,6</sup> | Sima Seifi<sup>6</sup> | Majid Ghayour-Mobarhan<sup>1,7</sup> |  
Gordon A. Ferns<sup>8</sup> | Amir Avan<sup>1,7</sup> | Seyed M. Hassanian<sup>1,6</sup>

<sup>1</sup>Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>2</sup>Department of Medical Genetics, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>3</sup>Department of Clinical Biochemistry and Immunogenetic Research Center, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Mazandaran, Iran

<sup>4</sup>Student Research Committee, Faculty of Medicine, Islamic Azad University, Mashhad, Iran

<sup>5</sup>Department of Nutrition, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>6</sup>Department of Medical Biochemistry, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>7</sup>Department of Modern Sciences and Technologies, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>8</sup>Brighton & Sussex Medical School, Division of Medical Education, Brighton, Sussex, UK

## Correspondence

Seyed M. Hassanian, Department of Medical Biochemistry, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.  
Email: hasanianmehrm@mums.ac.ir

## Funding information

Mashhad University of Medical Sciences, Grant/Award Numbers: 950852, 951824

## Abstract

The red blood cell distribution width (RDW) is an index of the heterogeneity of circulating red blood cell size, which along with other standard complete blood count (CBC) parameters are used to identify hematological system diseases. Besides hematological disorders, several clinical studies have shown that an increased in the RDW may be associated with other diseases including acute pancreatitis, chronic kidney disease, gastrointestinal disorders, cancer, and of special interest in this review, cardiovascular disease (CVD). The diagnostic and prognostic value of RDW in different CVD (acute coronary syndrome, ischemic cerebrovascular disease, peripheral artery disease, atrial fibrillation, heart failure, and acute ischemic stroke) has been reviewed in this article, to provide an understanding how its measurement may be applied to improve the management of these conditions.

## KEYWORDS

biomarker, cardiovascular diseases, RDW

**Abbreviations:** ACS, acute coronary syndrome; AF, atrial fibrillation; aSAH, aneurysmal subarachnoid hemorrhage; CABG, coronary artery by-pass graft; CBC, complete blood count; CNS, Canadian Neurological Scale; CRP, C-reactive protein; CVD, cardiovascular disease; GCS, Glasgow Coma Scale; HF, heart failure; hsCRP, high sensitivity C-reactive protein; IHD, ischemic heart disease; MCV, mean corpuscular volume; NIHSS, National Institutes of Health Stroke Scale; NSTEMI, non ST evaluation myocardial infraction; NT pro-BNP, N-terminal pro-brain natriuretic peptide; PAD, peripheral artery disease; PCI, percutaneous coronary intervention; RDW, red blood cell distribution width; SCAD, Stable coronary artery disease; SD, standard deviation; STEMI, ST elevation myocardial infraction; UA, unstable angina.

The authors have no financial conflicts of interest.

## 1 | INTRODUCTION

Cardiovascular disease (CVD) remains an important cause of mortality, and has a huge burden on health<sup>1</sup> especially in developing countries. Lifestyle and age are two key factors in the development of CVD.<sup>2</sup> The incidence of CVDs is increasing worldwide and comprises several specific conditions that include: ischemic heart disease (IHD), cerebrovascular disease, hypertensive heart disease, peripheral vascular disease, heart failure (HF), and atrial fibrillation (AF).<sup>3</sup> Ischemic heart disease and stroke are currently responsible for most cases of CVD.<sup>4</sup> There are multiple risk factors associated with CVD,<sup>4</sup> which can be divided into classical risk factors, that have been known for decades (including smoking, diabetes mellitus, hypertension, abnormal lipids, and obesity) and novel risk factors such as C-reactive protein (CRP), cystatin C, homocysteine, adipokines, adiponectin, resistin, osteoprotegerin, genetic factors, and red blood cell distribution width (RDW).<sup>1</sup> Most people at risk have several risk factors and multiple risk factors combination may have a synergistic impact to increase risk of CVD.<sup>4,5</sup>

The RDW is a simple, rapid, and easily available assessment measure of RBC size heterogeneity, which can be calculated from the standard deviation (SD) of erythrocyte volumes for the mean corpuscular volume (MCV).<sup>6</sup> The reference range variation is due to difference in the instruments used to measure the RDW.<sup>7</sup> However the common minimum and maximum of RDW physiological range are 11% and 15%, respectively.<sup>6</sup>

RDW is a routinely reported as a part of complete blood count. It has recently been reported that there is a possible relationship between RDW and many clinical conditions other than anemia. These conditions may alter RDW by various mechanisms that include: inflammation, nutritional disturbances, and alteration in erythropoiesis. Such alteration usually effect bone marrow and the hormone erythropoietin. This hormone regulates the production, maturation, and survival of erythrocytes. An abnormal increase in erythropoietin production will result in increase in RDW.<sup>8</sup> Inflammation inhibits erythropoietin and RBC production as well as reducing the survival of RBCs by damaging RBC memberane and increasing fragility because of increased oxidative stress.<sup>9</sup> Moreover, deficiencies of iron,<sup>10</sup> vitamin B12,<sup>11</sup> and folate<sup>12</sup> are considered as the main factors, which are directly related to the nutritional status. Other clinical conditions, which can increase RDW, are hemolytic anemia and sickle cell anemia are hematological diseases.<sup>7</sup> Moreover, an elevated RDW is also observed in non-hematological diseases that may interfere with red blood cell production because of alteration in overall inflammatory status. Such changes are seen with inflammatory bowel disease,<sup>13</sup> systemic lupus erythematosus,<sup>14</sup> hepatobiliary disease,<sup>15</sup> and Behçet's disease.<sup>7,16</sup>

There are several studies showing that an increased RDW is related to CVD complications.<sup>17,18</sup> RDW at the upper limit of normal range is associated with a significantly increased risk of

CVD and its mortality.<sup>19</sup> The exact mechanism behind the relationship of RDW values and developing different pathologies are not clearly understood. Anisocytosis can directly affect many clinical conditions including heart failure. High anisocytosis RBCs usually indicated the decreased oxygen capacity. Moreover, abnormal RBCs may participate in development of cardiac fibrosis by amplification of inflammation.<sup>8</sup> Such effects will result in decreased oxygenation of many organs including cardiomyocytes and predispose human cardiovascular system to different pathologies including ischemia and organ failure.<sup>20</sup> The purpose of this review was to summarize the results of recent studies that have investigated the diagnostic and prognostic value of RDW in CVD.

## 2 | RDW AS A BIOMARKER IN CORONARY ARTERY DISEASE

There are several studies investigating the association of increased RDW level with CVD complications. For instance, Lippi et al. assessed 456 acute coronary syndrome (ACS) and 1848 non-ACS patients and reported that median RDW level was significantly higher in ACS patient compared to non-ACS control group. Furthermore, it was shown that the sensitivity of ACS diagnosis could be improved by measuring cardiac troponin T and RDW simultaneously.<sup>21</sup> Uyarel et al. investigated the association between the RDW on admission with incidence of in-hospital mortality rate in 2506 patients with ST elevation myocardial infarction (STEMI) undergoing primary percutaneous coronary intervention (PCI). The authors showed that compared with patients with a normal RDW on admission, patients with increased RDW had higher in-hospital mortality rate.<sup>22</sup> Interestingly, investigation on 370 patients with STEMI and 156 normal coronary angiographic subjects showed that, in young patients, an increased RDW was associated with STEMI while this association was not observed in elderly patients.<sup>23</sup> Furthermore, the RDW was correlated with the number of stenotic coronary artery as well as fragmented QRS in patients with non-ST evaluation myocardial infraction (NSTEMI) or unstable angina (UA).<sup>24</sup> Similarly, it has been reported that a high RDW at the time of hospital admission in patients with stable coronary artery disease (SCAD) is associated with an enhanced mortality and incidence of ACS within the subsequent year.<sup>25</sup> RDW may also be helpful in evaluation of risk of mortality after ACS.<sup>26</sup> Nabais et al. studied 1,796 patients with ACS, and showed that increased RDW is an independent risk factor for 6-month mortality or MI in patients with ACS.<sup>27</sup> Poludasu et al. studied patients who underwent PCI and showed that an increased RDW was a risk factor for mortality in patients with normal level of hemoglobin at baseline.<sup>28</sup> An evaluation of 560 patients with diabetes mellitus and SCAD who underwent PCI showed that there was a significant association between RDW and long-term mortality in these patients.<sup>29</sup> Skjelbakken

et al. reported a positive association between RDW and risk of MI in the general population. The risk of MI was increased by ~ 13% for every 1% increment in RDW.<sup>30</sup> Gul et al. studied 310 patients with UA or NSTEMI and found that the RDW at time of hospital admission was a significant predictor of worse outcomes in these patients and mortality in the next 3 years was significantly higher in the patients with high RDW level.<sup>31</sup> Isik et al. followed up 193 patients who underwent coronary angiography and found that there was a higher RDW in patients who underwent angiography compared with those subjects with normal coronary arteries and coronary artery disease (CAD).<sup>32</sup> Dabbah et al. studied 1,709 patients with acute MI and results showed that RDW on admission and RDW elevation during the period of hospitalization were associated with poor clinical outcomes and mortality in patients with MI with and without anemia.<sup>33</sup> Vaya et al. studied the association between CVD events recurrence with RDW level in MI patients and found that high level of RDW increased the risk of CVD events by sixfold.<sup>34</sup> It has been consistently demonstrated that there is an association between high RDW level at the time of hospital admission with high sensitivity C-reactive protein (hsCRP), N-terminal pro-brain natriuretic peptide (NT pro-BNP), presence of more severe coronary arteries lesions and intracoronary thrombotic burden in patients with STEMI who underwent primary PCI.<sup>35</sup> In another study, 251 patients who had a bare-metal stent implantation, and who underwent control coronary angiography were investigated by Kurtul et al. to evaluate relationship between in-stent restenosis and level of RDW. They report that RDW level could predict in-stent restenosis before control coronary angiography.<sup>36</sup>

### 3 | PROGNOSTIC ROLE OF RDW IN ISCHEMIC CEREBROVASCULAR DISEASE

There are several studies investigating the association between RDW and stroke. Söderholm et al. measured RDW in 26,879 participants from the general population, without a history of previous stroke or CAD event, and found that subjects who had RDW level in the highest quartile had an increased subsequent incidence of stroke and cerebral infarction. However, no significant relationship was found between RDW and cerebral hemorrhage.<sup>37</sup> Furthermore, it has been shown that there is a positive correlation between both hs-CRP and RDW with carotid intimal-medial thickness in patients suffering from primary ischemic stroke. This study also suggested RDW may be a predictor of the presence carotid artery atherosclerosis.<sup>38</sup> Vayá et al. evaluated the correlation between RDW and stroke in 163 patients hospitalized due to cryptogenic stroke and 186 healthy subjects who were adjusted according to age, sex, anemia, lipid profile, and other potential confounders. They found that a high RDW was associated with the increase risk of


cryptogenic stroke.<sup>39</sup> Moreover, in a study on 847 patients with a first-ever episode of acute cerebral infarction it was found that high value of RDW was correlated with adverse functional outcome.<sup>40</sup> Ramírez-Moreno et al. measured the RDW in 224 patients with first-ever ischemic stroke and 224 cases without cerebrovascular disease and found a stepwise association between RDW levels and risk of cerebral infarction.<sup>41</sup> A higher RDW was also observed in 88 stroke patients with a poor Glasgow Coma Scale (GCS), National Institutes of Health Stroke Scale (NIHSS) as well as Canadian Neurological Scale (CNS). This study showed that the RDW may be a biomarker for the prediction of severity and functional outcomes in stroke patients.<sup>42</sup> Siegler et al. undertook a retrospectively investigated of RDW in 179 patients with aneurysmal subarachnoid hemorrhage (aSAH) admitted to a neurointensive care unit and found that there is a significant association between elevated RDW level and incidence of cerebral infarction and adverse clinical outcomes after aSAH.<sup>43</sup> Kaya et al. showed that in 153 patients with HF who had had a stroke there was a significantly increased baseline RDW in comparison with patients without stroke, suggesting that RDW may be a predictive marker for the incidence of stroke in HF patients. A RDW >15.2% at the time of admission had 87% sensitivity and 74% specificity for stroke incidence in HF patients.<sup>44</sup> Taken together, these results support the proposition that RDW could be a useful biomarker for stroke with a reasonable predictive value in these patients.

### 4 | ROLE OF RDW IN PERIPHERAL ARTERY DISEASE AND RDW

There is compelling evidence showing that RDW may be a biomarker of PAD. It has been consistently reported that a high RDW is associated with an increased risk of death in patients suffering from peripheral artery disease (PAD). In these patients, the risk of mortality increased almost 10% for every 1% increment in RDW.<sup>45</sup> Zalawadiya et al. measured RDW in 6,950 participants from general population and after adjustment for potential confounders they showed that, there is a direct relationship between RDW and prevalence of PAD.<sup>46</sup> Demirtas et al. studied 82 PAD patients and showed that RDW increased incrementally with the progression of PAD severity stages.<sup>47</sup>

### 5 | RDW A NOVEL BIOMARKER FOR ATRIAL FIBRILLATION

Several studies have evaluated the association between RDW and AF. Ertaş et al. studied 132 patients undergoing coronary artery by-pass graft (CABG) and measured AF before and after surgery in these patients. Preoperative RDW was significantly lower in patients who developed postoperative AF. However, no association was observed between the postoperative RDW



**FIGURE 1** The effects of increased RDW level on different types of CVD, supporting the clinical significance of this molecule in cardiovascular diseases. CVD, cardiovascular disease; RDW, red blood cell distribution width

**FIGURE 2** Association between increased RDW and CVD. CVD, cardiovascular disease; RDW, red blood cell distribution width

value and postoperative AF. Further studies showed that the sensitivity and specificity for prediction of postoperative AF at the level of RDW >13.4% were 61 and 60%, respectively.<sup>48</sup> Korantzopoulos et al. have reported that the sensitivity and specificity to predict incidence of AF at the preoperative level of RDW >13.3% were 80 and 60%, respectively, in patients undergoing cardiac surgery.<sup>49</sup> Similarly, it has been demonstrated that the RDW was significantly higher in patients with paroxysmal

non-valvular AF compared with patients without this. They also reported that RDW of >12.5% had 48% sensitivity and 67% specificity for the incidence of paroxysmal AF.<sup>50</sup> Güngör et al. showed patients with AF had a higher RDW compared to the control group, suggesting that increased RDW may be a risk marker for non-valvular AF.<sup>51</sup> Adamsson Eryd et al. followed up 27,124 participants without a prior history of AF for ~ 13 years, and reported that there was a positive association



**TABLE 1** The reported associations between RDW and CVDs

Author	Type of CVD	Study population	Main findings	Ref.
Lippi et al. (2009)	CAD	456 ACS patients and 1848 patients with non-ACS chest pain	RDW might be considered with other conventional cardiac markers for the risk stratification of ACS patients admitted to emergency departments.	12
Uyarel et al. (2011)	CAD	2,506 STEMI patients undergoing primary PCI	A high admission RDW level in patients with STEMI undergoing primary PCI was associated with increased risk for in-hospital and long-term cardiovascular mortality.	13
Uysal et al. (2012)	CAD	370 patients with STEMI and 156 subjects as control groups	High levels of RDW were associated with STEMI in young patients.	14
Bekler et al. (2014)	CAD	251 patients with NSTEMI or UA	An elevated RDW values is associated with fQRS in NST-ACS.	15
Ren et al. (2013)	CAD	1,442 patients with SCAD	An elevated RDW value is associated with an increased risk of 1-year adverse outcomes in patients with stable angina pectoris.	16
Timóteo et al. (2015)	CAD	787 patients with ACS	Combining both RDW and PDW markers with GRACE risk score improved the predictive value for all-cause mortality and reduced the estimated risk of those who did not die.	17
Nabais et al. (2009)	CAD	1,796 patients with ACS	A graded independent was found association between higher RDW values and adverse outcomes in patients with ACS.	18
Poludasu et al. (2009)	CAD	859 patients who underwent PCI	Higher RDW is a strong and independent predictor of long-term mortality in patients undergoing PCI who were not anemic at baseline.	19
Tsuboi et al. (2013)	CAD	560 diabetic patients with SCAD underwent elective PCI	Increased RDW was significantly associated with increased long-term all-cause mortality in diabetic patients after PCI.	8
Skjelbakken et al. (2014)	CAD	25,612 participants from general population	RDW is associated with incident MI in a general population independent of anemia and cardiovascular risk factors.	20
Gul et al. (2012)	CAD	310 patients with NSTEMI or UA	RDW is a readily available clinical laboratory value associated with long-term cardiovascular mortality in NSTEMI and UAP.	21
Isik et al. (2012)	CAD	193 patients who underwent coronary angiography for SCAD	RDW is a readily available clinical laboratory value that is associated with both the presence and the complexity of CAD.	22
Dabbah et al. (2010)	CAD	1,709 patients with MI	There is a graded, independent association between increased RDW and mortality after AMI.	23
Vaya et al. (2012)	CAD	119 patients with MI	RDW seems to be an independent predictor for recurrent cardiovascular events in AMI patients.	24
Wang et al. (2015)	CAD	424 patients with STEMI who underwent primary PCI	A high RDW may be associated with the severity and instability of acute myocardial infarction.	25
Kurtul et al. (2015)	CAD	251 patients with a bare-metal stent underwent control coronary angiography	Increased serum RDW levels were independently associated with bare-metal ISR in patients with stable coronary artery disease.	26

(Continues)

TABLE 1 (Continued)

Author	Type of CVD	Study population	Main findings	Ref.
Söderholm et al. (2015)	ICD	26,879 participants from general population without a history of coronary events or stroke	Red cell distribution width in the highest quartile was associated with increased incidence of total stroke and cerebral infarction.	27
Jia et al. (2015)	ICD	392 patients with primary ischemic stroke	RDW level were higher than non-CAS in patients with primary ischemic stroke.	28
Vayá et al. (2015)	ICD	163 patients with cryptogenic stroke and 186 healthy controls	The RDW >14% increased the risk of CS by 2.5-fold, irrespectively of anemia, inflammation, and lipidic profile.	29
Kim et al. (2012)	ICD	847 patients with a first-ever episode of acute cerebral infarction	Higher RDW measured in cases of acute stage cerebral infarction was associated with poor functional outcome and mortality.	30
Ramírez-Moreno et al. (2013)	ICD	224 patients with first-ever ischemic stroke	Based on this study, RDW is a powerful predictor of stroke.	31
Kara et al. (2015)	ICD	88 patients with acute ischemic stroke	In stroke patients who have symptoms <24 hr, the RDW may be useful in predicting the severity and functional outcomes of the stroke.	32
Siegler et al. (2017)	ICD	179 patients with aSAH	RDW elevation is associated with cerebral infarction and poor outcome after aSAH.	33
Kaya et al. (2015)	ICD	153 HF patients	RDW may be important hematological indices for stroke in patients with HF using propensity score analysis.	34
Ye et al. (2011)	PAD	13,039 patients with PAD	RDW, a routinely available measurement, is an independent prognostic marker in patients with PAD.	35
Zalawadiya et al. (2012)	PAD	6,950 participants from general population	Higher levels of RDW are independently associated with a higher risk of PAD and can significantly improve the risk prediction beyond that estimated by ACC/AHA-defined PAD screening criteria.	36
Demirtas et al. (2014)	PAD	82 patients with PAD	Both red blood cell distribution width and mean platelet volume are found to be associated with the severity of atherosclerotic disease in patients with peripheral arterial disease.	37
Ertas et al. (2013)	AF	132 patients with post-operative AF undergoing CABG	Preoperative RDW level predicts new-onset AF after CABG in patients without histories of AF.	38
Güngör et al. (2014)	AF	117 patients with AF and 60 controls	Elevated RDW levels, not NLR, may be an independent risk marker for non-valvular AF.	41
Adamsson et al. (2014)	AF	27,124 participants with no history of AF, HF, MI, or stroke, followed up for a mean of 13 years	Red blood cell distribution width was associated with incidence of AF independently of several cardiovascular, nutritional, and hematological factors in this study of middle-aged subjects from the general population.	42
Gurses et al. (2015)	AF	299 patients with AF followed for a median period of 24 months	Elevated RDW may be a predictor of late recurrence following cryoballoon-based AF ablation.	43
Wan et al. (2015)	AF	300 patients with AF followed up for a median up period of 3.2 years	Elevated RDW is an independent predictor of long-term adverse clinical outcomes, including all-cause mortality and MAEs, among patients with AF.	44

(Continues)

**TABLE 1** (Continued)

Author	Type of CVD	Study population	Main findings	Ref.
Kaya et al. (2017)	AF	619 patients with AF	Increased level of RDW is independently correlated with higher risk for development of LAS in patients with non-valvular AF.	45
Emans et al. (2013)	HF	17,533 participants from general population followed up for a mean period of 11-years	RDW is associated with HF events in an apparently healthy middle-aged population.	46
Borné et al. (2011)	HF	26,784 subjects without history of MI, stroke or HF, followed up for a mean period of 15 years	RDW was found to be associated with long-term incidence of first hospitalization due to HF among middle-aged subjects.	47
Oh et al. (2009)	HF	100 patients with acute HF	A novel relation was found between higher levels of RDW and elevated E/E' in patients with AHF.	48
Nishizaki et al. (2012)	HF	160 patients died in the Department of Cardiology	HF was the major cause of death, and RDW $\geq 16.5$ was significantly associated with fatal HF in super-elderly patients.	49
Uemura et al. (2016)	HF	229 patients with acutely decompensated HF	The change in RDW values during hospitalization independently predicts poor outcomes in patients with ADHF.	50
Huang et al. (2014)	HF	18,288 patients with HF	HF patients with higher RDW may have poorer prognosis than those with lower RDW.	51
Xanthopoulos et al. (2017)	HF	218 patients with both HF and DM	RDW at admission was associated with higher event rate both in HF patients with and without DM.	52
Turcato et al. (2017)	HF	588 ADHF patients	More aggressive management may be advisable in ADHF patients with increasing anisocytosis during the first days of hospitalization.	53
Bilal et al. (2016)	HHD	100 hypertensive patients	A significant number of patients with hypertension have increased levels of RDW.	54
Ozcan et al. (2013)	HHD	247 patient with essential hypertension	RDW, as easy and quick measurable tool, can predict non-dipping pattern in essential hypertension.	55
Fici et al. (2013)	HHD	72 essential hypertensive patients	Nebivolol improved RDW and N/L to a greater extent than metoprolol in patients with hypertension.	57

Abbreviations: ACS, acute coronary syndrome; ADHF, acutely decompensated HF; AF, atrial fibrillation; aSAH, aneurysmal subarachnoid hemorrhage; CABG, coronary artery by-pass graft; CAD, coronary artery disease; DM, diabetes mellitus; HF, heart failure; HHD, Hypertensive heart disease; hs-CRP, high sensitivity C-reactive protein; ICD, Ischemic cerebrovascular disease; IHD, ischemic heart disease; MI, myocardial infarction; NSTEMI, non-ST elevation myocardial infarction; OSAHS, obstructive sleep apnea hypopnea syndrome; PAD, Peripheral Artery Disease; PCI, percutaneous coronary intervention; RDW, red blood cell distribution width; SCAD, stable coronary artery disease; STEMI, ST elevation myocardial infarction; UA, unstable angina.

between RDW and risk of AF in the study group.<sup>52</sup> Moreover, RDW was shown to be associated with recurrence of late AF after monitoring 299 patients with AF for a mean of 2 years.<sup>53</sup> To further investigate whether elevated RDW is associated with the mortality rate in AF patients, Wan et al. followed up AF patients for a mean of 3 years and found that increased RDW level was associated with an enhanced mortality rate, and with major adverse events in these patients.<sup>54</sup> Kaya et al. studied 619 patients with non-valvular AF and showed that increased level of RDW was associated with both left atrial stasis and thromboembolic events in patients with non-valvular AF.<sup>55</sup>

These results clearly suggest that RDW could be a valid biomarker for both prognosis and diagnosis in AF patients.

## 6 | PROGNOSTIC AND DIAGNOSTIC VALUE OF RDW IN HEART FAILURE

In recent years growing evidence suggest that RDW is a key player in increasing risk of incidence and poor prognosis of heart failure (HF). Emans et al. studied 17,533 subjects for a mean period of 11-years and found a nonlinear association

between RDW level and risk of HF even after adjusting for other risk factors including C-reactive protein, ferritin and iron. They also reported that there was an inverse association between RDW level and physical inactivity, however this association was not correlated with HF.<sup>56</sup> Borné et al. studied 26,784 healthy subjects with no prior history of stroke, MI, or HF, and followed up them for a mean of 15 years to investigate the association between RDW and long-term incidence of hospitalization for HF. Patients with a high RDW had significantly higher risk of HF incidence, but after adjustment for other risk factors including hs-CRP, N-terminal pro-B-type natriuretic peptide and cystatin C, the significance of the association between RDW and HF incidence was reduced substantially.<sup>57</sup> Furthermore, an elevated RDW was shown to be correlated with the increased left ventricular filling pressure in patients with acute HF.<sup>58</sup> To further support the clinical significance of RDW in HF patients, Nishizaki et al. evaluated 160 patients who died as a result of HF and found that RDW >16.5% was significantly associated with fatal HF.<sup>59</sup> Uemura et al. measured RDW at the time of hospital admission and during hospitalization in patients with acute decompensated HF and reported that the alteration in RDW level during hospital stay, can predict adverse outcomes and all-cause death among patients.<sup>60</sup> The results of a meta-analysis of 18,288 HF patients showed that level of RDW on admission and discharge, as well as its variation during treatment, can be used as marker to predict prognosis of HF. This meta-analysis also revealed that HF patients with higher RDW had poorer prognosis.<sup>61</sup> The RDW can be high in patients who have both HF and diabetes mellitus (DM). Xanthopoulos et al. studied 218 patients with both HF and DM and measured RDW at hospital admission, discharge and after 4, 8, 12 months of discharge. There was an association between RDW on admission and higher incidence of clinical events in diabetic or non-diabetic HF. However, it was shown that RDW 1 year after discharge was significantly higher in HF patients with DM compared to non-diabetic HF patients.<sup>62</sup> It has been demonstrated that increased RDW after 4 days of hospitalization, increased risk of 30-day mortality up to fourfold in patients with acute decompensated HF, suggesting that these patients with elevated RDW require more care in the first days of hospitalization.<sup>63</sup>

## 7 | HYPERTENSIVE HEART DISEASE AND RDW

Several studies suggested that RDW level is increased in hypertensive patients. Tanindi et al. showed that RDW is increased in prehypertensive and hypertensive patients compared with healthy subjects independent of other factors such as age and inflammatory status. Moreover, it has been demonstrated that elevated RDW is strongly associated with systolic and diastolic blood pressures.<sup>18</sup> Bilal et al. evaluated 100 hypertensive

patients and found that most of these patients had an increased level of RDW.<sup>64</sup> Similarly, Ozcan et al. validated the association between RDW and hypertension in 247 patients with essential hypertension and found that RDW is significantly higher in patients with non-dipper hypertension compared with the dipper hypertensive patients.<sup>65</sup> Furthermore, it has been demonstrated that nebivolol, an anti-hypertensive drug, decreased RDW in new essential hypertensive patients, supporting the hypothesis that RDW could be used in clinic as a novel biomarker for evaluating the hypertension-associated complications in cardiovascular disease.<sup>66</sup>

## 8 | CONCLUSION

Several studies have investigated the relationship of RDW with CVD (Table 1). RDW either alone or in combination with other cardiac biomarkers may improve diagnosis and assessing prognosis of CVD including ACS, ischemic cerebrovascular disease, peripheral artery disease, atrial fibrillation, heart failure, and acute ischemic stroke. In brief, elevated RDW has unfavorable effects on CVD and is associated with increased risk of incidence, adverse outcome, and mortality of CVD. The effects of increased RDW on different type of CVD are showed in Figure 1.

The exact relevant mechanisms are unclear and not yet fully understood (Figure 2). Although RDW may be a biomarker that is rapid and easily available assessment in almost all health facilities and obtained as a part of complete blood count, for evaluating the severity and prognosis of patients with CVDs, further investigations are needed to assess the efficacy as well as accuracy of RDW in CVDs.

## ORCID

Seyed M. Parizadeh  <https://orcid.org/0000-0001-5543-0436>

Majid Ghayour-Mobarhan  <https://orcid.org/0000-0002-1081-6754>

Amir Avan  <https://orcid.org/0000-0002-4968-0962>

Seyed M. Hassanian  <https://orcid.org/0000-0002-5247-4043>

## REFERENCES

1. Payne RA. Cardiovascular risk. *Br J Clin Pharmacol*. 2012;74(3):396–410.
2. Dahlöf B. Cardiovascular disease risk factors: Epidemiology and risk assessment. *Am J Cardiol*. 2010;105(1 Suppl):3A–9A.
3. Bowry A, Lewey J, Dugani S, Choudhry N. The burden of cardiovascular disease in low-and middle-income countries: Epidemiology and management. *Can J Cardiol*. 2015;31(9):1151–1159.
4. Norhammar A, Schenck-Gustafsson K. Type 2 diabetes and cardiovascular disease in women. *Diabetologia*. 2013;56(1):1–9.





5. Tran D, Zimmerman L. Cardiovascular risk factors in young adults: A literature review. *J Cardiovasc Nurs*. 2015;30(4):298–310.
6. Danese E, Lippi G, Montagnana M. Red blood cell distribution width and cardiovascular diseases. *J Thorac Dis*. 2015;7(10):E402–E411.
7. Zöller B, Melander O, Svensson P, Engström G. Red cell distribution width for predicting cardiovascular disease: A literature review. *EMJ Cardiol*. 2014;2:61–70.
8. Salvagno GL, Sanchis-Gomar F, Picanza A, Lippi G. Red blood cell distribution width: A simple parameter with multiple clinical applications. *Crit Rev Clin Lab Sci*. 2015;52(2):86–105.
9. Feng G-H, Li H-P, Li Q-L, Fu Y, Huang R-B. Red blood cell distribution width and ischaemic stroke. *Stroke Vasc Neurol*. 2017;2(3):172–175.
10. Sultana GS, Haque SA, Sultana T, Ahmed AN. Value of red cell distribution width (RDW) and RBC indices in the detection of iron deficiency anemia. *Mymensingh Med J*. 2013;22(2):370–376.
11. Saxena S, Weiner JM, Carmel R. Red blood cell distribution width in untreated pernicious anemia. *Am J Clin Pathol*. 1988;89(5):660–663.
12. Dugdale AE. Predicting iron and folate deficiency anaemias from standard blood testing: The mechanism and implications for clinical medicine and public health in developing countries. *Theor Biol Med Model*. 2006;3(34):1742–4682.
13. Oliveira AM, Cardoso FS, Rodrigues CG, et al. Can red cell distribution width be used as a marker of Crohn's disease activity? *GE Port J Gastroenterol*. 2015;23(1):6–12.
14. Zou XL, Lin XJ, Ni X, Wang J, Liu W, Wei J. Baseline red blood cell distribution width correlates with disease activity and therapeutic outcomes in patients with systemic lupus Erythematosus, irrespective of anemia status. *Clin Lab*. 2016;62(10):1841–1850.
15. Hu Z-D. Red blood cell distribution width: A promising index for estimating activity of autoimmune disease. *J Lab Precis Med*. 2016;1(2):62–68.
16. Shafiee M, Tayefi M, Hassanian SM, et al. Depression and anxiety symptoms are associated with white blood cell count and red cell distribution width: A sex-stratified analysis in a population-based study. *Psychoneuroendocrinology*. 2017;84:101–108.
17. Isik T, Ayhan E, Kurt M, Tanboga IH, Kaya A, Aksakal E. Is red cell distribution width a marker for the presence and poor prognosis of cardiovascular disease? *Eur J Med*. 2012;44(3):169–171.
18. Tanindi A, Topal F, Topal F, Celik B. Red cell distribution width in patients with prehypertension and hypertension. *Blood Press*. 2012;21(3):177–181.
19. Su C, Liao L-Z, Song Y, Xu Z-W, Mei W-Y. The role of red blood cell distribution width in mortality and cardiovascular risk among patients with coronary artery diseases: A systematic review and meta-analysis. *J Thorac Dis*. 2014;6(10):1429–1440.
20. Lippi G, Turcato G, Cervellin G, Sanchis-Gomar F. Red blood cell distribution width in heart failure: A narrative review. *World J Cardiol*. 2018;10(2):6–14.
21. Lippi G, Filippozzi L, Montagnana M, et al. Clinical usefulness of measuring red blood cell distribution width on admission in patients with acute coronary syndromes. *Clin Chem Lab Med*. 2009;47(3):353–357.
22. Uyarel H, Ergelen M, Cicek G, et al. Red cell distribution width as a novel prognostic marker in patients undergoing primary angioplasty for acute myocardial infarction. *Coron Artery Dis*. 2011;22(3):138–144.
23. Uysal O, Duran M, Ozkan B, et al. Red cell distribution width is associated with acute myocardial infarction in young patients. *Cardiol J*. 2012;19(6):597–602.
24. Bekler A, Gazi E, Tenekecioglu E, et al. Assessment of the relationship between red cell distribution width and fragmented QRS in patients with non-ST elevated acute coronary syndrome. *Med Sci Monit*. 2014;20:413.
25. Ren H, Hua Q, Quan M, et al. Relationship between the red cell distribution width and the one-year outcomes in Chinese patients with stable angina pectoris. *Inter Med (Tokyo, Japan)*. 2013;52(16):1769–1774.
26. Timóteo A, Papoila A, Lousinha A, et al. Predictive impact on medium-term mortality of hematological parameters in acute coronary syndromes: Added value on top of GRACE risk score. *Eur Heart J Acute Cardiovasc Care*. 2015;4(2):172–179.
27. Nabais S, Losa N, Gaspar A, et al. Association between red blood cell distribution width and outcomes at six months in patients with acute coronary syndromes. *Rev Port Cardiol*. 2009;28(9):905.
28. Poludasu S, Marmur J, Weedon J, Khan W, Cavusoglu E. Red cell distribution width (RDW) as a predictor of long-term mortality in patients undergoing percutaneous coronary intervention. *Thromb Haemost*. 2009;102(3):581–587.
29. Tsuboi S, Miyauchi K, Kasai T, et al. Impact of red blood cell distribution width on long-term mortality in diabetic patients after percutaneous coronary intervention. *Jpn Circ J*. 2013;77(2):456–461.
30. Skjelbakken T, Lappegård J, Ellingsen TS, et al. Red cell distribution width is associated with incident myocardial infarction in a general population: The Tromsø study. *J Am Heart Assoc*. 2014;3(4):83–92.
31. Gul M, Uyarel H, Ergelen M, et al. The relationship between red blood cell distribution width and the clinical outcomes in non-ST elevation myocardial infarction and unstable angina pectoris: A 3-year follow-up. *Coron Artery Dis*. 2012;23(5):330–336.
32. Isik T, Uyarel H, Tanboga I, et al. Relation of red cell distribution width with the presence, severity, and complexity of coronary artery disease. *Coron Artery Dis*. 2012;23(1):51–56.
33. Dabbah S, Hammerman H, Markiewicz W, Aronson D. Relation between red cell distribution width and clinical outcomes after acute myocardial infarction. *Am J Cardiol*. 2010;105(3):312–317.
34. Vaya A, Hernández J, Zorio E, Bautista D. Association between red blood cell distribution width and the risk of future cardiovascular events. *Clin Hemorheol Microcirc*. 2012;50(3):221.
35. Wang P, Wang Y, Li H, Wu Y, Chen H. Relationship between the red blood cell distribution width and risk of acute myocardial infarction. *J Atheroscler Thromb*. 2015;22(1):21–26.
36. Kurtul A, Murat S, Yarlioglu M, et al. The association of red cell distribution width with in-stent restenosis in patients with stable coronary artery disease. *Platelets*. 2015;26(1):48–52.
37. Söderholm M, Borné Y, Hedblad B, Persson M, Engström G. Red cell distribution width in relation to incidence of stroke and carotid atherosclerosis: A population-based cohort study. *PLoS ONE*. 2015;10(5):e0124957.
38. Jia H, Li H, Zhang Y, Li C, Hu Y, Xia C. Association between red blood cell distribution width (RDW) and carotid artery atherosclerosis (CAS) in patients with primary ischemic stroke. *Arch Gerontol Geriatr*. 2015;61(1):72–75.
39. Vayá A, Hernández V, Rivera L, et al. Red blood cell distribution width in patients with cryptogenic stroke. *Clin Appl Thromb Hemost*. 2015;21(3):241–245.

40. Kim J, Kim Y, Song T, et al. Red blood cell distribution width is associated with poor clinical outcome in acute cerebral infarction. *Thromb Haemost.* 2012;108(2):349–356.
41. Ramírez-Moreno J, Gonzalez-Gomez M, Ollero-Ortiz A, Roa-Montero A, Gómez-Baquero M, Constantino-Silva A. Relation between red blood cell distribution width and ischemic stroke: A case-control study. *Int J Stroke.* 2013;8(6):E36.
42. Kara H, Degirmenci S, Bayir A, et al. Red cell distribution width and neurological scoring systems in acute stroke patients. *Neuropsychiatr Dis Treat.* 2015;11:733.
43. Siegler J, Marcaccio C, Nawalinski K, et al. Elevated red cell distribution width is associated with cerebral infarction in aneurysmal subarachnoid hemorrhage. *Neurocrit Care.* 2017;26(1):26–33.
44. Kaya A, Isik T, Kaya Y, et al. Relationship between red cell distribution width and stroke in patients with stable chronic heart failure: A propensity score matching analysis. *Clin Appl Thromb Hemost.* 2015;21(2):160–165.
45. Ye Z, Smith C, Kullo IJ. Usefulness of red cell distribution width to predict mortality in patients with peripheral artery disease. *Am J Cardiol.* 2011;107(8):1241–1245.
46. Zalawadiya S, Veeranna V, Panaich S, Afonso L. Red cell distribution width and risk of peripheral artery disease: Analysis of National Health and nutrition examination survey 1999–2004. *Vasc Med (London, England).* 2012;17(3):155–163.
47. Demirtas S, Karahan O, Yazici S, et al. The relationship between complete blood count parameters and Fontaine's stages in patients with peripheral arterial disease. *Vascular.* 2014;22(6):427–431.
48. Ertaş G, Aydın C, Sönmez O, et al. Red cell distribution width predicts new-onset atrial fibrillation after coronary artery bypass grafting. *Scand Cardiovasc J.* 2013;47(3):132–135.
49. Korantzopoulos P, Liu T. RDW as a marker of postoperative atrial fibrillation. *Int J Cardiol.* 2015;191:109.
50. Liu T, Shao Q, Miao S, et al. Red cell distribution width as a novel, inexpensive marker for paroxysmal atrial fibrillation. *Int J Cardiol.* 2014;171(2):e52–e53.
51. Güngör B, Özcan K, Erdinler İ, et al. Elevated levels of RDW is associated with non-valvular atrial fibrillation. *J Thromb Thrombolysis.* 2014;37(4):404–410.
52. Adamsson ES, Borné Y, Melander O, et al. Red blood cell distribution width is associated with incidence of atrial fibrillation. *J Intern Med.* 2014;275(1):84–92.
53. Gurses K, Yalcin M, Kocyigit D, et al. Red blood cell distribution width predicts outcome of cryoballoon-based atrial fibrillation ablation. *J Interv Card Electrophysiol.* 2015;42(1):51–58.
54. Wan H, Yang Y, Zhu J, et al. The relationship between elevated red cell distribution width and long-term outcomes among patients with atrial fibrillation. *Clin Biochem.* 2015;48(12):762–767.
55. Kaya A, Tukkan C, Alper AT, et al. Increased levels of red cell distribution width is correlated with presence of left atrial stasis in patients with non-valvular atrial fibrillation. *North Clin Istanbul.* 2017;4(1):66–72.
56. Emans M, Gaillard C, Pfister R, et al. Red cell distribution width is associated with physical inactivity and heart failure, independent of established risk factors, inflammation or iron metabolism; the EPIC-Norfolk study. *Int J Cardiol.* 2013;168(4):3550–3555.
57. Borné Y, Smith J, Melander O, Hedblad B, Engström G. Red cell distribution width and risk for first hospitalization due to heart failure: A population-based cohort study. *Eur J Heart Fail.* 2011;13(12):1355–1361.
58. Oh J, Kang S, Hong N, et al. Relation between red cell distribution width with echocardiographic parameters in patients with acute heart failure. *J Card Fail.* 2009;15(6):517–522.
59. Nishizaki Y, Yamagami S, Suzuki H, et al. Red blood cell distribution width as an effective tool for detecting fatal heart failure in super-elderly patients. *Inter Med (Tokyo, Japan).* 2012;51(17):2271–2276.
60. Uemura Y, Shibata R, Takemoto K, et al. Elevation of red blood cell distribution width during hospitalization predicts mortality in patients with acute decompensated heart failure. *J Cardiol.* 2016;67(3):268–273.
61. Huang Y-L, Hu Z-D, Liu S-J, et al. Prognostic value of red blood cell distribution width for patients with heart failure: A systematic review and meta-analysis of cohort studies. *PLoS ONE.* 2014;9(8):120–127.
62. Xanthopoulos A, Giamouzis G, Melidonis A, et al. Red blood cell distribution width as a prognostic marker in patients with heart failure and diabetes mellitus. *Cardiovasc Diabetol.* 2017;16:81.
63. Turcato G, Zorzi E, Prati D, et al. Early in-hospital variation of red blood cell distribution width predicts mortality in patients with acute heart failure. *Int J Cardiol.* 2017;243:306–310.
64. Bilal A, Farooq JH, Kiani I, Assad S, Ghazanfar H, Ahmed I. Importance of mean red cell distribution width in hypertensive patients. *Cureus.* 2016;8(11):902–909.
65. Ozcan F, Turak O, Durak A, et al. Red cell distribution width and inflammation in patients with non-dipper hypertension. *Blood Press.* 2013;22(2):80–85.
66. Fici F, Celik T, Balta S, et al. Comparative effects of nebivolol and metoprolol on red cell distribution width and neutrophil/lymphocyte ratio in patients with newly diagnosed essential hypertension. *J Cardiovasc Pharmacol.* 2013;62(4):388–393.

**How to cite this article:** Parizadeh SM, Jafarzadeh-Esfehani R, Bahreyni A, et al. The diagnostic and prognostic value of red cell distribution width in cardiovascular disease; current status and prospective. *BioFactors.* 2019;1–10. <https://doi.org/10.1002/biof.1518>